

For patients with **ALK+ Metastatic Non-Small Cell Lung Cancer**



# Your Guide to Treatment With **ALUNBRIG** (brigatinib)

This guide  
**is designed**  
**to help** you  
get started  
and stay on  
track with  
**ALUNBRIG.**

ACTOR PORTRAYAL

## What Is **ALUNBRIG**?

ALUNBRIG is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC):

- that has a certain type of abnormal anaplastic lymphoma kinase (ALK) gene, **and**
- that has spread to other parts of your body


It is not known if ALUNBRIG is safe and effective in children.

**Please read the Important Safety Information throughout and the [Patient Information](#) in the accompanying [full Prescribing Information](#).**

# How to take **ALUNBRIG** (brigatinib)

**FOLLOW YOUR HEALTHCARE PROVIDER'S INSTRUCTIONS.** ALUNBRIG is a tablet that is taken orally once a day. Your healthcare provider will start you on a 90 mg dose of ALUNBRIG for the first 7 days of treatment.

If you tolerate this dose of ALUNBRIG well, after the first 7 days of treatment, your healthcare provider may increase your dose to 180 mg orally once daily. Be sure to review your ALUNBRIG dosing regimen with your healthcare team to ensure you are taking it correctly.



**BE SURE TO REVIEW YOUR ALUNBRIG DOSING REGIMEN** with your healthcare team to ensure you are taking it correctly. Do not stop taking ALUNBRIG without talking to your healthcare provider.

**Please read the Important Safety Information throughout and the Patient Information in the accompanying [full Prescribing Information](#).**

## **One tablet**, once a day, with or without food



### **ONE TABLET**

comes in convenient 180-mg, 90-mg, and 30-mg tablets



### **ONCE DAILY**

take it at the same time every day



### **WITH OR WITHOUT FOOD**

no need to plan meals around the medication—eat whenever and whatever you want



### **AVOID GRAPEFRUIT**

and grapefruit juice because it may change the amount of ALUNBRIG in your blood



### **STORE ALUNBRIG**

at room temperature of 68°F to 77°F (20°C to 25°C)

## **THINGS TO REMEMBER**

- Take ALUNBRIG exactly as your healthcare provider tells you to take it. Do not change your dose or stop taking ALUNBRIG unless your healthcare provider tells you to
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with ALUNBRIG if you develop side effects
- Swallow ALUNBRIG tablets whole. Do not crush or chew tablets
- If you miss a dose of ALUNBRIG, do not take the missed dose. Take your next dose at your regular time
- If you vomit after taking a dose of ALUNBRIG, do not take an extra dose. Take your next dose at your regular time

Use the calendar to remind yourself to take **ALUNBRIG** (brigatinib) on schedule and track your dose

Write down your dose for each day as prescribed by your healthcare provider, and then check the box when you have taken that dose.

Try to take **ALUNBRIG** at the same time every day.



	Week ONE	Week TWO	Week THREE	Week FOUR
DAY 1	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 2	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 3	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 4	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 5	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 6	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 7	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____

## Tips for taking your medication daily

**Taking your medication every day is important. Consider these tips to help you stay on track and ensure that you are taking your daily dose of ALUNBRIG as scheduled.**

- Take your medication every day around the same time, such as when you have breakfast. Keep your pill box or medication bottle on or near the breakfast table
- Set an alarm in your house or on your mobile device to remind you to take your medication
- Designate a paper or online calendar to track your daily dose and mark each day with an X after you've taken your medication
- Ask a friend, family member, or caregiver to remind you to take your medication
- Be sure to take your medication with you when you travel. If you are traveling by air, keep your medication in your carry-on bag. If you do leave it at home, call your healthcare provider right away. Your healthcare team may be able to get a new supply of ALUNBRIG to you at your new location
- If you are struggling with side effects, contact your healthcare provider right away to discuss the side effects you are experiencing. Your healthcare team may be able to help you manage them

**AFTER WEEK 4**, continue to take your medication at the same dose unless your healthcare provider changes your treatment plan. If you need to refill your prescription, contact your healthcare provider or pharmacist.

**Please read the Important Safety Information throughout and the Patient Information in the accompanying full Prescribing Information.**

# ALUNBRIG (brigatinib) can cause other serious side effects

## WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW

**ABOUT ALUNBRIG? ALUNBRIG can cause serious side effects, including:**

**Lung problems. ALUNBRIG may cause severe or life-threatening swelling (inflammation) of the lungs any time during treatment, and can lead to death.** These lung problems happen especially within the first week of treatment with ALUNBRIG. Symptoms may be similar to those symptoms from lung cancer. Tell your healthcare provider right away if you have any new or worsening symptoms, including: trouble breathing or shortness of breath, chest pain, cough (with or without mucus), or fever.

**High blood pressure (hypertension).** ALUNBRIG may cause high blood pressure. Your healthcare provider will check your blood pressure before starting and during treatment with ALUNBRIG. Tell your healthcare provider right away if you get headaches, dizziness, blurred vision, chest pain, or shortness of breath.

**Slow heart rate (bradycardia).** ALUNBRIG may cause very slow heartbeats that can be severe. Your healthcare provider will check your heart rate during treatment with ALUNBRIG. Tell your healthcare provider right away if you feel dizzy, lightheaded, or faint during treatment with ALUNBRIG. Tell your healthcare provider if you start to take

or have any changes in heart rate or blood pressure medicines.

**Vision problems.** ALUNBRIG may cause vision problems. Your healthcare provider may stop ALUNBRIG and refer you to an eye specialist if you develop severe vision problems during treatment with ALUNBRIG. Tell your healthcare provider right away if you have any loss of vision or any change in vision, including double vision, seeing flashes of light, blurry vision, light hurting your eyes, or new or increased floaters.

**Muscle pain, tenderness, and weakness (myalgia).** ALUNBRIG may increase the level of an enzyme in your blood called creatine phosphokinase (CPK), which may be a sign of muscle damage. Your healthcare provider will do blood tests to check your blood levels of CPK during treatment with ALUNBRIG. Tell your healthcare provider right away if you get new or worsening signs and symptoms of muscle problems, including unexplained muscle pain or muscle pain that does not go away, tenderness, or weakness.

**Inflammation of the pancreas (pancreatitis).** ALUNBRIG may increase enzymes in your blood called amylase and lipase, which may be a sign of pancreatitis. Your healthcare provider will do blood tests to check your pancreatic enzyme blood levels during treatment with ALUNBRIG. Tell your healthcare provider right away if you

get new or worsening signs and symptoms of pancreatitis, including upper abdominal pain that may spread to the back and get worse with eating, weight loss, or nausea.

**High blood sugar (hyperglycemia).** ALUNBRIG may increase your blood sugar levels. Your healthcare provider will do blood tests to check your blood sugar levels before starting and during treatment with ALUNBRIG. Your healthcare provider may need to start or change your blood sugar medicine to control your blood sugar levels. Tell your healthcare provider right away if you get new or worsening signs and symptoms of hyperglycemia, including needing to urinate more than usual, or feeling very thirsty, very hungry, sick to your stomach, weak, tired, or confused.

**Please read the Important Safety Information throughout and the Patient Information in the accompanying full Prescribing Information.**



# Additional Important Safety Information for ALUNBRIG (brigatinib)

The most common side effects of ALUNBRIG include diarrhea, fatigue, nausea, rash, cough, muscle pain, headache, high blood pressure, vomiting and difficulty breathing.

ALUNBRIG may cause fertility problems in males. This may affect your ability to father a child. Talk to your healthcare provider if you have concerns about fertility.

These are not all of the possible side effects of ALUNBRIG.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Before you take ALUNBRIG, tell your healthcare provider about all of your medical conditions, including if you have lung or breathing problems, high blood pressure, a slow heartbeat, or any vision problems, have or have had pancreatitis, have diabetes mellitus or glucose intolerance, have liver problems, have kidney problems or are on dialysis, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

ALUNBRIG can harm your unborn baby. Your healthcare provider will determine whether or not you are pregnant before you start treatment with ALUNBRIG. Tell your healthcare provider right away if you become pregnant during treatment with ALUNBRIG or think you may be pregnant.

• **Females** who are able to become pregnant should use effective non-hormonal birth control during treatment with ALUNBRIG and for at least 4 months after the final dose of ALUNBRIG. Birth control pills (oral contraceptives) and other hormonal forms of birth control may not be effective if used during treatment with ALUNBRIG. Talk to your healthcare provider about birth control choices that are right for you during treatment with ALUNBRIG.

• **Males** who have female partners that are able to become pregnant should use effective birth control during treatment with ALUNBRIG and for at least 3 months after the final dose of ALUNBRIG.

It is not known if ALUNBRIG passes into your breast milk. Do not breastfeed during treatment with ALUNBRIG and for 1 week after the final dose of ALUNBRIG.

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, or herbal supplements.

Avoid eating grapefruit or drinking grapefruit juice during treatment with ALUNBRIG. Grapefruit may increase the amount of ALUNBRIG in your blood.



Please read the [Patient Information](#) in the accompanying [full Prescribing Information](#).

# Side effect tracker

DATE	TIME		SIGNS OR SYMPTOMS OF SIDE EFFECTS
	START	END	

SHARE YOUR TRACKER with your healthcare team

Please read the Important Safety Information throughout and the Patient Information in the accompanying full Prescribing Information.

# Communicate with your healthcare team regularly

**BE SURE TO TELL THEM ABOUT ANY SIDE EFFECTS YOU MAY EXPERIENCE**

- **ALUNBRIG (brigatinib) can cause** side effects, some may be serious
- **Talk to your healthcare provider** if any side effects occur
- **Keep track of how you are feeling** and record any side effects you may be experiencing
- **Share your side effect tracker with your healthcare provider** and be sure to discuss any new or worsening side effects right away
- **Tell your healthcare provider about all of the medicines you take** because some medicines may affect the way ALUNBRIG works

## IMPORTANT NUMBERS

Fill in the contact information for your healthcare team and keep this list on hand.

### ONCOLOGIST

NAME

PHONE

EMAIL

### AFTER HOURS CONTACT

NAME

PHONE

EMAIL

### PRIMARY CARE HEALTHCARE PROVIDER

NAME

PHONE

EMAIL

### AFTER HOURS CONTACT

NAME

PHONE

EMAIL

### PHARMACIST

NAME

PHONE

EMAIL

### AFTER HOURS CONTACT

NAME

PHONE

EMAIL

Please read the Important Safety Information throughout and the [Patient Information](#) in the accompanying [full Prescribing Information](#).



ACTOR PORTRAYAL

## Patient Support Program



From finding financial assistance that may be right for you to understanding your disease, Takeda Oncology Here2Assist can provide the information you need throughout your treatment.

Takeda Oncology Here2Assist:

- Works with your healthcare provider and your insurance company to help get you started on your medication
- Identifies available financial assistance that may be right for you
- May help get you started on treatment if there is a delay in insurance coverage determination
- Works with your pharmacy to arrange the delivery of your medication
- Connects you to additional support services and resources

For more information on Takeda Oncology Here2Assist, call 1-844-817-6468, Option 2 or visit [www.Here2Assist.com](http://www.Here2Assist.com).

**LET'S TALK. We're available Monday-Friday, 8 AM-8 PM ET.  
Call 1-844-817-6468, Option 2 for information on  
Takeda Oncology Here2Assist.**



ONCOLOGY

All trademarks are the property of their respective owners. ©2020 Millennium Pharmaceuticals, Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Limited. All rights reserved. 06/20 MAT-USO-BRG-19-00046