

# MEET ERIC

## A REAL PATIENT LIVING WITH ALK+ mNSCLC

### Medical, Family, and Social History:

- Supply chain professional
- Leads a busy and active lifestyle, which includes golfing, traveling, working, and spending time with friends and family
- Nonsmoker
- No known family history of cancer



## LEARN ABOUT ERIC'S TREATMENT JOURNEY INSIDE.

### INDICATION

ALUNBRIG® (brigatinib) is indicated for the treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.

### IMPORTANT SAFETY INFORMATION WARNINGS AND PRECAUTIONS

#### Interstitial Lung Disease (ILD)/Pneumonitis

Severe, life-threatening, and fatal pulmonary adverse reactions consistent with interstitial lung disease (ILD)/pneumonitis have occurred with ALUNBRIG. In ALTA 1L, ILD/pneumonitis occurred in 5.1% of patients receiving ALUNBRIG. ILD/pneumonitis occurred within 8 days of initiation of ALUNBRIG in 2.9% of patients, with Grade 3 to 4 reactions occurring in 2.2% of patients. In ALTA, ILD/pneumonitis occurred in 3.7% of patients in the 90 mg group (90 mg once daily) and 9.1% of patients in the 90→180 mg group (180 mg once daily with 7-day lead-in at 90 mg once daily). Adverse reactions consistent with possible ILD/pneumonitis occurred within 9 days of initiation of ALUNBRIG (median onset was 2 days) in 6.4% of patients, with Grade 3 to 4 reactions occurring in 2.7% of patients. Monitor for new or worsening respiratory symptoms (dyspnea, cough, etc.), particularly during the first week of initiating ALUNBRIG. Withhold ALUNBRIG in any patient with new or worsening respiratory symptoms, and promptly evaluate for ILD/pneumonitis or other causes of respiratory symptoms (e.g., pulmonary embolism, tumor progression, and infectious pneumonia). For Grade 1 or 2 ILD/pneumonitis, either resume ALUNBRIG with dose reduction according to Table 1 of the full Prescribing Information after recovery to baseline or permanently discontinue ALUNBRIG. Permanently discontinue ALUNBRIG for Grade 3 or 4 ILD/pneumonitis or recurrence of Grade 1 or 2 ILD/pneumonitis.

ALK, anaplastic lymphoma kinase; mNSCLC, metastatic non-small cell lung cancer.

**Please see additional Important Safety Information throughout and accompanying full Prescribing Information.**



## ERIC'S DIAGNOSIS AND TREATMENT JOURNEY

### Diagnosis of ALK+ mNSCLC:

- In late 2021, Eric had a bad cough that he attributed to a respiratory virus. However, following worsening of his cough and the development of back pain and dyspnea, Eric had a virtual checkup before visiting his primary care physician
- Following an initial CT chest scan, Eric received a needle-guided biopsy and additional testing, including X-ray, CT, and PET scans, that suggested he had NSCLC
- NGS testing was conducted, and metastatic ALK rearrangement was confirmed
- MRI results confirmed presence of brain metastases
- 3 weeks after his initial checkup, Eric, aged 41 at the time, was diagnosed with ALK+ mNSCLC

### Prior to Initiating Treatment, Eric's CT Scans Revealed a Tumor in the Left Lung and Additional Metastases

- Eric's CT scans revealed left lung consolidation and right lung nodules
- Enlarged lymph nodes and pericardial effusion were also observed
- Eric's MRI revealed brain metastases at baseline

## HOW WOULD YOU TREAT A PATIENT LIKE ERIC?

CT, computerized tomography; MRI, magnetic resonance imaging; PET, positron emission tomography.

### WARNINGS AND PRECAUTIONS (continued)

#### Hypertension

In ALTA 1L, hypertension was reported in 32% of patients receiving ALUNBRIG; 13% of patients experienced Grade 3 hypertension. In ALTA, hypertension was reported in 11% of patients in the 90 mg group and 21% of patients in the 90→180 mg group. Grade 3 hypertension occurred in 5.9% of patients overall. Control blood pressure prior to treatment with ALUNBRIG. Monitor blood pressure after 2 weeks and at least monthly thereafter during treatment with ALUNBRIG. Withhold ALUNBRIG for Grade 3 hypertension despite optimal antihypertensive therapy. Upon resolution or improvement to Grade 1, resume ALUNBRIG at the same dose. Consider permanent discontinuation of treatment with ALUNBRIG for Grade 4 hypertension or recurrence of Grade 3 hypertension. Use caution when administering ALUNBRIG in combination with antihypertensive agents that cause bradycardia.

2 Please see additional Important Safety Information throughout and accompanying full Prescribing Information.

## ERIC'S 1L TREATMENT GOALS

- An option with strong 1L systemic efficacy
- Due to his brain metastases, he prefers a treatment that can offer strong intracranial efficacy in the first line
- Continue to travel with his wife and spend time with family
- Maintain his quality of life with an option that is well tolerated

### ALUNBRIG (brigatinib) Was Initiated at 90 mg Once Daily for the First 7 Days, Then the Dose was Increased to 180 mg Once Daily<sup>1</sup>

“

*When choosing my first-line treatment, we evaluated the multiple efficacious options with my doctors and determined that **the intracranial efficacy of ALUNBRIG, as well as the side effect profile, made it the best choice for me.***

– Eric

”



## SEE ERIC'S RESULTS ON THE BACK COVER.

1L, first line.

### WARNINGS AND PRECAUTIONS (continued)

#### Bradycardia

In ALTA 1L, heart rates less than 50 beats per minute (bpm) occurred in 8.1% of patients receiving ALUNBRIG; one patient (0.7%) experienced Grade 3 bradycardia. In ALTA, heart rates less than 50 beats per minute (bpm) occurred in 5.7% of patients in the 90 mg group and 7.6% of patients in the 90→180 mg group. One patient (0.9%) in the 90 mg group experienced Grade 2 bradycardia. Monitor heart rate and blood pressure during treatment with ALUNBRIG. Monitor patients more frequently if concomitant use of drug known to cause bradycardia cannot be avoided. For symptomatic bradycardia, withhold ALUNBRIG and review concomitant medications for those known to cause bradycardia. If a concomitant medication known to cause bradycardia is identified and discontinued or dose adjusted, resume ALUNBRIG at the same dose following resolution of symptomatic bradycardia; otherwise, reduce the dose of ALUNBRIG following resolution of symptomatic bradycardia. Discontinue ALUNBRIG for life-threatening bradycardia if no contributing concomitant medication is identified.

  
ALUNBRIG®  
BRIGATINIB  
180mg | 90mg | 30mg  
TABLETS

## IMPORTANT SAFETY INFORMATION (continued)

### WARNINGS AND PRECAUTIONS (continued)

#### Visual Disturbance

In ALTA 1L, Grade 1 or 2 adverse reactions leading to visual disturbance, including blurred vision, photophobia, photopsia, and reduced visual acuity, were reported in 7.4% of patients receiving ALUNBRIG. In ALTA, adverse reactions leading to visual disturbance, including blurred vision, diplopia, and reduced visual acuity, were reported in 7.3% of patients treated with ALUNBRIG in the 90 mg group and 10% of patients in the 90→180 mg group. Grade 3 macular edema and cataract occurred in one patient each in the 90→180 mg group. Advise patients to report any visual symptoms. Withhold ALUNBRIG and obtain an ophthalmologic evaluation in patients with new or worsening visual symptoms of Grade 2 or greater severity. Upon recovery of Grade 2 or Grade 3 visual disturbances to Grade 1 severity or baseline, resume ALUNBRIG at a reduced dose. Permanently discontinue treatment with ALUNBRIG for Grade 4 visual disturbances.

#### Creatine Phosphokinase (CPK) Elevation

In ALTA 1L, creatine phosphokinase (CPK) elevation occurred in 81% of patients who received ALUNBRIG. The incidence of Grade 3 or 4 CPK elevation was 24%. Dose reduction for CPK elevation occurred in 15% of patients. In ALTA, CPK elevation occurred in 27% of patients receiving ALUNBRIG in the 90 mg group and 48% of patients in the 90→180 mg group. The incidence of Grade 3 to 4 CPK elevation was 2.8% in the 90 mg group and 12% in the 90→180 mg group. Dose reduction for CPK elevation occurred in 1.8% of patients in the 90 mg group and 4.5% of patients in the 90→180 mg group. Advise patients to report any unexplained muscle pain, tenderness, or weakness. Monitor CPK levels during ALUNBRIG treatment. Withhold ALUNBRIG for Grade 3 or 4 CPK elevation with Grade 2 or higher muscle pain or weakness. Upon resolution or recovery to Grade 1 CPK elevation or baseline, resume ALUNBRIG at the same dose or at a reduced dose per Table 2 of the full Prescribing Information.

#### Pancreatic Enzyme Elevation

In ALTA 1L, amylase elevation occurred in 52% of patients and Grade 3 or 4 amylase elevation occurred in 6.8% of patients who received ALUNBRIG. Lipase elevations occurred in 59% of patients and Grade 3 or 4 lipase elevation occurred in 17% of patients. In ALTA, amylase elevation occurred in 27% of patients in the 90 mg group and 39% of patients in the 90→180 mg group. Lipase elevations occurred in 21% of patients in the 90 mg group and 45% of patients in the 90→180 mg group. Grade 3 or 4 amylase elevation occurred in 3.7% of patients in the 90 mg group and 2.7% of patients in the 90→180 mg group. Grade 3 or 4 lipase elevation occurred in 4.6% of patients in the 90 mg group and 5.5% of patients in the 90→180 mg group. Monitor lipase and amylase during treatment with ALUNBRIG. Withhold ALUNBRIG for Grade 3 or 4 pancreatic enzyme elevation. Upon resolution or recovery to Grade 1 or baseline, resume ALUNBRIG at the same dose or at a reduced dose.

#### Hepatotoxicity

In ALTA 1L, aspartate aminotransferase (AST) elevations occurred in 72% of patients and Grade 3 or 4 AST elevations occurred in 4.5% of patients who received ALUNBRIG. Alanine aminotransferase (ALT) elevations occurred in 52% of patients and Grade 3 or 4 ALT elevations occurred in 5.2% of patients. One patient (0.7%) had a serious adverse reaction of hepatocellular injury. In ALTA, AST elevations occurred in 38% of patients in the 90 mg group and 65% of patients in the 90→180 mg group. ALT elevations occurred in 34% of patients in the 90 mg group and 40% of patients in the 90→180 mg group. Grade 3 or 4 AST elevations occurred in 0.9% of patients in the 90 mg group and did not occur in any patients in the 90→180 mg group. Grade 3 or 4 ALT elevations did not occur in any patients in the 90 mg group and in 2.7% of patients in the 90→180 mg group. Monitor AST, ALT and total bilirubin during treatment with ALUNBRIG, especially during the first 3 months. Withhold ALUNBRIG for Grade 3 or 4 hepatic enzyme elevation with bilirubin less than or equal to 2 × ULN. Upon resolution or recovery to Grade 1 or less (less than or equal to 3 × ULN) or to baseline, resume ALUNBRIG at a next lower dose per Table 2 of the full Prescribing Information. Permanently discontinue ALUNBRIG for Grade 2 to 4 hepatic enzyme elevation with concurrent total bilirubin elevation greater than 2 times the ULN in the absence of cholestasis or hemolysis.

## WARNINGS AND PRECAUTIONS (continued)

#### Hyperglycemia

In ALTA 1L, 56% of patients who received ALUNBRIG experienced new or worsening hyperglycemia. Grade 3 hyperglycemia, based on laboratory assessment of serum fasting glucose levels, occurred in 7.5% of patients. In ALTA, 43% of patients who received ALUNBRIG experienced new or worsening hyperglycemia. Grade 3 hyperglycemia, based on laboratory assessment of serum fasting glucose levels, occurred in 3.7% of patients. Two of 20 (10%) patients with diabetes or glucose intolerance at baseline required initiation of insulin while receiving ALUNBRIG. Assess fasting serum glucose prior to initiation of ALUNBRIG and monitor periodically thereafter. Initiate or optimize anti-hyperglycemic medications as needed. If adequate hyperglycemic control cannot be achieved with optimal medical management, withhold ALUNBRIG until adequate hyperglycemic control is achieved and consider reducing the dose of ALUNBRIG dosage per Table 1 of the full Prescribing Information or permanently discontinuing ALUNBRIG.

#### Photosensitivity

In ALTA 1L, 3.7% of patients who received ALUNBRIG experienced photosensitivity, with 0.7% of patients experiencing Grade 3 to 4 reactions. In ALTA, 0.9% of patients who received ALUNBRIG in the 90 mg group and 0.9% of patients in the 90→180 mg group experienced photosensitivity. Grade 3 to 4 photosensitivity was not reported in patients in the 90 mg group or in the 90→180 mg group. Advise patients to limit sun exposure while taking ALUNBRIG, and for at least 5 days after discontinuation of treatment. Advise patients, when outdoors, to wear a hat and protective clothing, and use a broad-spectrum Ultraviolet A (UVA)/Ultraviolet B (UVB) sunscreen and lip balm (SPF ≥30) to help protect against sunburn. Based on the severity, withhold ALUNBRIG, then resume at the same dose, or reduce the dose, or permanently discontinue.

#### Embryo-Fetal Toxicity

Based on its mechanism of action and findings in animals, ALUNBRIG can cause fetal harm when administered to pregnant women. There are no clinical data on the use of ALUNBRIG in pregnant women. Advise women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with ALUNBRIG and for at least 4 months following the final dose. Advise males with female partners of reproductive potential to use effective contraception during treatment and for at least 3 months after the last dose of ALUNBRIG.

### ADVERSE REACTIONS

The most common adverse reactions (≥25%) with ALUNBRIG were diarrhea, fatigue, nausea, rash, cough, myalgia, headache, hypertension, vomiting, and dyspnea.

### DRUG INTERACTIONS

**CYP3A Inhibitors:** Avoid coadministration of ALUNBRIG with strong or moderate CYP3A inhibitors. If coadministration of a strong or moderate CYP3A inhibitor is unavoidable, reduce the dose of ALUNBRIG.

**CYP3A Inducers:** Avoid coadministration of ALUNBRIG with strong or moderate CYP3A inducers. If coadministration of a moderate CYP3A inducer is unavoidable, increase the dose of ALUNBRIG.

# ERIC EXPERIENCED STRONG RESULTS WITH 1L ALUNBRIG (brigatinib)

3 Months After His ALUNBRIG Initiation,

## Eric Achieved a Response in Both the Lung and Brain

- He achieved 90% reduction of tumor volume in lung
  - Left lung consolidation significantly improved; right lung nodular metastases had also improved
  - Mild improvement of pericardial effusion
  - No enlarged lymph nodes in the chest, abdomen, or pelvis
- Achieved a 50% reduction of tumor volume in the brain
  - Interval resolution of perilesional edema
  - Multiple intracranial lesions have decreased in size
  - Additional lesions previously described within the frontal and parietal lobes were no longer visualized, with no new enhancing intracranial lesions detected

## Adverse Events Experienced by Eric:

- Bradycardia for first 3 months, which was monitored at home until it naturally resolved
- Nausea and stomach pain managed with prescription anti-nausea medication and ginger lozenges
- No dose reductions occurred

*The most common adverse reactions (≥25%) with ALUNBRIG were diarrhea, fatigue, nausea, rash, cough, myalgia, headache, hypertension, vomiting, and dyspnea seen in patients other than Eric.*

## As of April 2024, Eric has:

- Remained on 1L ALUNBRIG since January 2022
- Retained his active lifestyle and quality of life

This is Eric's experience with ALUNBRIG. Individual results may vary.

**WILL YOU CONSIDER ALUNBRIG FOR YOUR NEXT PATIENT LIKE ERIC?**

## USE IN SPECIFIC POPULATIONS

### Females and Males of Reproductive Potential

Verify pregnancy status in females of reproductive potential prior to initiating ALUNBRIG. Advise females of reproductive potential to use effective contraception during treatment with ALUNBRIG and for at least 4 months after the final dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with ALUNBRIG and for at least 3 months after the final dose. ALUNBRIG may cause reduced fertility in males.

**Lactation:** Advise patients not to breastfeed.

**Hepatic Impairment:** Reduce the dose of ALUNBRIG for patients with severe hepatic impairment.



**Renal Impairment:** Reduce the dose of ALUNBRIG for patients with severe renal impairment.

To report SUSPECTED ADVERSE REACTIONS, contact Takeda Pharmaceuticals U.S.A., Inc. at 1-844-217-6468 or the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**Please see additional Important Safety Information throughout and accompanying full Prescribing Information.**



ONCOLOGY

TAKEDA and  are registered trademarks of Takeda Pharmaceutical Company Limited. ALUNBRIG and  are registered trademarks of ARIAD Pharmaceuticals, Inc.  
© 2024 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 09/24 US0-BRG-0843

