



Your Guide to Treatment With **ALUNBRIG**[®] (brigatinib)

This guide
is designed
to help you
get started
and stay on
track with
ALUNBRIG.

ACTOR PORTRAYAL

What is ALUNBRIG?

ALUNBRIG is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC):

- that has a certain type of abnormal anaplastic lymphoma kinase (ALK) gene, **and**
- that has spread to other parts of your body


It is not known if ALUNBRIG is safe and effective in children.

Please read the Important Safety Information throughout and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

How to take ALUNBRIG (brigatinib)

FOLLOW YOUR HEALTHCARE PROVIDER'S INSTRUCTIONS. ALUNBRIG is a tablet that is taken orally once a day. Your healthcare provider will start you on a 90 mg dose of ALUNBRIG for the first 7 days of treatment.

If you tolerate this dose of ALUNBRIG well, after the first 7 days of treatment, your healthcare provider may increase your dose to 180 mg orally once daily. Be sure to review your ALUNBRIG dosing regimen with your healthcare team to ensure you are taking it correctly.



BE SURE TO REVIEW YOUR ALUNBRIG DOSING REGIMEN with your healthcare team to ensure you are taking it correctly. Do not stop taking ALUNBRIG without talking to your healthcare provider.

Please read the Important Safety Information throughout and the Patient Information in the accompanying full Prescribing Information.

One tablet, once a day, with or without food



ONE TABLET

comes in convenient 180-mg, 90-mg, and 30-mg tablets



ONCE DAILY

take it at the same time every day



WITH OR WITHOUT FOOD

no need to plan meals around the medication—eat whenever and whatever you want



AVOID GRAPEFRUIT

and grapefruit juice because it may change the amount of ALUNBRIG in your blood



STORE ALUNBRIG

at room temperature of 68°F to 77°F (20°C to 25°C)

THINGS TO REMEMBER

- Take ALUNBRIG exactly as your healthcare provider tells you to take it. Do not change your dose or stop taking ALUNBRIG unless your healthcare provider tells you to
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with ALUNBRIG if you develop side effects
- Swallow ALUNBRIG tablets whole. Do not crush or chew tablets
- If you miss a dose of ALUNBRIG, do not take the missed dose. Take your next dose at your regular time
- If you vomit after taking a dose of ALUNBRIG, do not take an extra dose. Take your next dose at your regular time

Use the calendar to remind yourself to take **ALUNBRIG** (brigatinib) on schedule and track your dose

Write down your dose for each day as prescribed by your healthcare provider, and then check the box when you have taken that dose.

Try to take ALUNBRIG at the same time every day.



	Week ONE	Week TWO	Week THREE	Week FOUR
DAY 1	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 2	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 3	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 4	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 5	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 6	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____
DAY 7	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____	Taken <input type="checkbox"/> _____ mg Time: _____

Tips for taking your medication daily

Taking your medication every day is important. Consider these tips to help you stay on track and ensure that you are taking your daily dose of ALUNBRIG as scheduled.

- Take your medication every day around the same time, such as when you have breakfast. Keep your pill box or medication bottle on or near the breakfast table
- Set an alarm in your house or on your mobile device to remind you to take your medication
- Designate a paper or online calendar to track your daily dose and mark each day with an X after you've taken your medication
- Ask a friend, family member, or caregiver to remind you to take your medication
- Be sure to take your medication with you when you travel. If you are traveling by air, keep your medication in your carry-on bag. If you do leave it at home, call your healthcare provider right away. Your healthcare team may be able to get a new supply of ALUNBRIG to you at your new location
- If you are struggling with side effects, contact your healthcare provider right away to discuss the side effects you are experiencing. Your healthcare team may be able to help you manage them

AFTER WEEK 4, continue to take your medication at the same dose unless your healthcare provider changes your treatment plan. If you need to refill your prescription, contact your healthcare provider or pharmacist.

Please read the Important Safety Information throughout and the Patient Information in the accompanying full Prescribing Information.

ALUNBRIG® (brigatinib) IMPORTANT SAFETY INFORMATION

What is ALUNBRIG?

ALUNBRIG is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC):

- that has a certain type of abnormal anaplastic lymphoma kinase (ALK) gene, **and**
- that has spread to other parts of your body

It is not known if ALUNBRIG is safe and effective in children.

ALUNBRIG can cause serious side effects, including:

- **Lung problems. ALUNBRIG may cause severe or life-threatening swelling (inflammation) of the lungs any time during treatment and can lead to death.** These lung problems happen **especially within the first week of treatment** with ALUNBRIG. Symptoms may be similar to those symptoms from lung cancer. Tell your healthcare provider right away if you have any new or worsening symptoms, including:
 - trouble breathing or shortness of breath
 - chest pain
 - cough with or without mucus
 - fever
- **High blood pressure (hypertension).** ALUNBRIG may cause high blood pressure. Your healthcare provider will check your blood pressure before starting and during treatment with ALUNBRIG. Tell your healthcare provider right away if you get headaches, dizziness, blurred vision, chest pain or shortness of breath.
- **Slow heart rate (bradycardia).** ALUNBRIG may cause very slow heartbeats that can be severe. Your healthcare provider will check your heart rate during treatment with ALUNBRIG. Tell your healthcare provider right away if you feel dizzy, lightheaded, or faint during treatment with ALUNBRIG. Tell your healthcare provider if you take any heart or blood pressure medicines.
- **Vision problems.** ALUNBRIG may cause vision problems. Your healthcare provider may stop ALUNBRIG and refer you to an eye specialist if you develop severe vision problems during treatment with ALUNBRIG. Tell your healthcare provider right away if you have any loss of vision or any change in vision, including:
 - double vision
 - seeing flashes of light
 - blurry vision
 - light hurting your eyes
 - new or increased floaters
- **Muscle pain, tenderness, and weakness (myalgia).** ALUNBRIG may increase the level of an enzyme in your blood called creatine phosphokinase (CPK), which may be a sign of muscle damage. Your healthcare provider will do blood tests to check your blood levels of CPK during treatment with ALUNBRIG. Tell your healthcare provider right away if you get new or worsening signs and symptoms of muscle problems, including unexplained muscle pain or muscle pain that does not go away, tenderness, or weakness.
- **Inflammation of the pancreas (pancreatitis).** ALUNBRIG may increase enzymes in your blood called amylase and lipase, which may be a sign of pancreatitis. Your healthcare provider will do blood tests to check your pancreatic enzyme blood levels during treatment with ALUNBRIG. Tell your healthcare provider right away if you get new or worsening signs and symptoms of pancreatitis, including upper abdominal pain that may spread to the back and get worse with eating, weight loss, or nausea.
- **Liver problems (hepatotoxicity).** ALUNBRIG may increase the levels of bilirubin in your blood and enzymes called aspartate aminotransferase (AST) and alanine aminotransferase (ALT) in your blood, which may be a sign of liver problems. Your healthcare provider will do blood tests to check your liver during treatment with ALUNBRIG. Tell your healthcare provider right away if you get new or worsening signs or symptoms, including:
 - yellowing of your skin or the white part of your eyes
 - dark or brown (tea color) urine
 - nausea or vomiting
 - pain on the right side of your stomach area
 - bleed or bruise more easily than normal
 - itchy skin
 - decreased appetite
 - feeling tired

Please read the Important Safety Information throughout and the Patient Information in the accompanying full [Prescribing Information](#).



• **High blood sugar (hyperglycemia).** ALUNBRIG may increase your blood sugar levels. Your healthcare provider will do blood tests to check your blood sugar levels before starting and during treatment with ALUNBRIG. Your healthcare provider may need to start or change your blood sugar medicine to control your blood sugar levels. Tell your healthcare provider right away if you get new or worsening signs and symptoms of hyperglycemia, including:

- feeling very thirsty
- feeling very hungry
- feeling weak or tired
- needing to urinate more than usual
- feeling sick to your stomach
- feeling confused

The most common side effects of ALUNBRIG include diarrhea, fatigue, nausea, rash, cough, muscle pain, headache, high blood pressure, vomiting and difficulty breathing.

ALUNBRIG may cause fertility problems in males. This may affect your ability to father a child. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of ALUNBRIG. Call your doctor for medical advice about side effects. [You may report side effects to the FDA at 1-800-FDA-1088.](#)

What should I avoid while taking ALUNBRIG?

- Limit your time in the sun during treatment with ALUNBRIG and for at least 5 days after your final dose. ALUNBRIG may make your skin sensitive to sunlight. You may burn more easily and get severe sunburns. When you are in the sun, wear a hat and protective clothing, and use a broad-spectrum sunscreen and lip balm with a Sun Protection Factor (SPF) of 30 or greater to protect against sunburn.
- Avoid eating grapefruit or drinking grapefruit juice during treatment with ALUNBRIG. Grapefruit may increase the amount of ALUNBRIG in your blood.

Before you take ALUNBRIG, tell your healthcare provider about all of your medical conditions, including if you have lung or breathing problems, high blood pressure, a slow heartbeat, or any vision problems, have or have had pancreatitis, have liver problems, have diabetes mellitus or glucose intolerance, have kidney problems or are on dialysis, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

- ALUNBRIG can harm your unborn baby. Your healthcare provider will determine whether or not you are pregnant before you start treatment with ALUNBRIG. Tell your healthcare provider right away if you become pregnant during treatment with ALUNBRIG or think you may be pregnant.
- **Females** who are able to become pregnant should use effective birth control during treatment with ALUNBRIG and for at least 4 months after the final dose of ALUNBRIG. Talk to your healthcare provider about birth control choices that are right for you during treatment with ALUNBRIG.
- **Males** who have female partners who are able to become pregnant should use effective birth control during treatment with ALUNBRIG and for at least 3 months after the final dose of ALUNBRIG.
 - It is not known if ALUNBRIG passes into your breast milk. Do not breastfeed during treatment with ALUNBRIG and for 1 week after the final dose of ALUNBRIG.

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, or herbal supplements.



Please read the [Patient Information](#) in the accompanying full [Prescribing Information](#) and discuss with your healthcare provider.

Communicate with your healthcare team regularly

BE SURE TO TELL THEM ABOUT ANY SIDE EFFECTS YOU MAY EXPERIENCE

- **ALUNBRIG (brigatinib) can cause** side effects, some may be serious
- **Talk to your healthcare provider** if any side effects occur
- **Keep track of how you are feeling** and record any side effects you may be experiencing
- **Share your side effect tracker with your healthcare provider** and be sure to discuss any new or worsening side effects right away
- **Tell your healthcare provider about all of the medicines you take** because some medicines may affect the way ALUNBRIG works

IMPORTANT NUMBERS

Fill in the contact information for your healthcare team and keep this list on hand.

ONCOLOGIST

NAME

PHONE

EMAIL

AFTER HOURS CONTACT

NAME

PHONE

EMAIL

PRIMARY CARE HEALTHCARE PROVIDER

NAME

PHONE

EMAIL

AFTER HOURS CONTACT

NAME

PHONE

EMAIL

PHARMACIST

NAME

PHONE

EMAIL

AFTER HOURS CONTACT

NAME

PHONE

EMAIL

Please read the Important Safety Information throughout and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

Patient Support Programs



From finding financial assistance that may be right for you to understanding your disease, Takeda Oncology Here2Assist™ can provide the information you need throughout your treatment.

Takeda Oncology Here2Assist:

- ▶ Works with your insurance company to help you get started on your medication
- ▶ Identifies available financial assistance that may be right for you
- ▶ May help get you started on treatment if there is a delay in insurance coverage determination
- ▶ Connects you to additional support services and resources
- ▶ Identifies specialty pharmacies to help fill and ship your prescriptions appropriately
- ▶ Conducts regular follow-up calls with you
- ▶ Sends you status updates and reminders via text messages*

Our case managers are your connection to personalized support



ACCESS SUPPORT: Once you're enrolled, Takeda Oncology Here2Assist case managers can work with you and your healthcare provider to determine your coverage options, and provide additional support throughout your treatment.



FINANCIAL ASSISTANCE: If you need assistance affording your medication, Takeda Oncology Here2Assist can help identify financial assistance programs that may be able to help you with the cost of your treatment.



HELPFUL RESOURCES: Takeda Oncology Here2Assist Program case managers can provide information about additional resources that may help address day-to-day concerns associated with your treatment.

To learn more about Takeda Oncology Here2Assist, call to speak with a case manager at 1-844-817-6468, Option 2, or visit www.Here2Assist.com. **Let's Talk.** We're available Monday-Friday, 8AM-8PM ET.

*Patients will need to enroll in the texting program to receive text messages.



ONCOLOGY

All trademarks are the property of their respective owners. ©2022 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved.
03/22 US0-BRG-0473